arXiv:1403.3827v1 [math.GR] 15 Mar 2014

CLASSIFYING ORBITS OF THE AFFINE GROUP OVER THE INTEGERS

LEONARDO MANUEL CABRER AND DANIELE MUNDICI

Abstract. For each $n = 1, 2, \ldots$, let $\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$ be the affine group over the integers. For every point $x = (x_1, \ldots, x_n) \in \mathbb{R}^n$ let $\mathrm{orb}(x) = \{\gamma(x) \in \mathbb{R}^n \mid \gamma \in \mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n\}$. Let G_x be the subgroup of the additive group $\mathbb{R}$ generated by $x_1, \ldots, x_n, 1$. If $\mathrm{rank}(G_x) \neq n$ then $\mathrm{orb}(x) = \{y \in \mathbb{R}^n \mid G_y = G_x\}$. Thus, G_x is a complete classifier of $\mathrm{orb}(x)$. By contrast, if $\mathrm{rank}(G_x) = n$, knowledge of G_x alone is not sufficient in general to uniquely recover $\mathrm{orb}(x)$: as a matter of fact, G_x determines precisely $\max(1, \frac{\phi(d)}{2})$ different orbits, where d is the denominator of the smallest positive nonzero rational in G_x, and ϕ is Euler function. To get a complete classification, rational polyhedral geometry provides an integer $1 \leq c_x \leq \max(1, d/2)$ such that $\mathrm{orb}(y) = \mathrm{orb}(x)$ iff $(G_x, c_x) = (G_y, c_y)$.

1. Introduction

Throughout we let $\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$ denote the group of affine transformations of the form $x \mapsto \mathcal{U}x + t$ for $x \in \mathbb{R}^n$, where $t \in \mathbb{Z}^n$ and $\mathcal{U}$ is an integer $n \times n$ matrix with $\det(\mathcal{U}) = \pm 1$. The dimension of the ambient space $\mathbb{R}^n$ will always be clear from the context. We let $\mathrm{orb}(x)$ denote the $(\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n)$-*orbit* of $x \in \mathbb{R}^n$,

$$\mathrm{orb}(x) = \{\gamma(x) \in \mathbb{R}^n \mid \gamma \in \mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n\}.$$

In this self-contained paper, generalizing [7] we classify $(\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n)$-orbits for all n: every point in $\mathbb{R}^n$ is assigned an invariant, in such a way that two points have the same $(\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n)$-orbit iff they have the same invariant. For each $x = (x_1, \ldots, x_n) \in \mathbb{R}^n$, the main invariant considered in this paper is the group

$$G_x = \mathbb{Z} + \mathbb{Z}x_1 + \mathbb{Z}x_2 + \cdots + \mathbb{Z}x_n$$

generated by $1, x_1, x_2, \ldots, x_n$ in the additive group $\mathbb{R}$. From Corollaries 16 and 17 it follows that G_x completely classifies the $(\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n)$-orbit of any $x \in \mathbb{R}^n$ when $\mathrm{rank}(G_x) \neq n$.

If $\mathrm{rank}(G_x) = n$, G_x determines precisely $\max(1, \frac{\phi(d)}{2})$ different $(\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n)$-orbits, where d is the denominator of the smallest positive nonzero rational in G_x and ϕ is Euler function. To get a complete invariant for $\mathrm{orb}(x)$, rational polyhedral geometry equips x with an integer $1 \leq c_x \leq \max(1, d/2)$ which, together with G_x, completely classifies $\mathrm{orb}(x)$. Thus in Theorem 15 we prove: $\mathrm{orb}(x) = \mathrm{orb}(y)$ iff $(G_x, c_x) = (G_y, c_y)$. In case $\mathrm{rank}(G_x) \neq n$, $c_x = 1$.

While the study of the orbits of the groups $\mathsf{GL}(2, \mathbb{Z})$, $\mathsf{SL}(2, \mathbb{Z})$, $\mathsf{SL}(2, \mathbb{Z})_+$ requires techniques from various mathematical areas [1, 2, 4, 8, 9, 10], a main possible reason of interest in our classification stems from the pervasive and novel role played by rational polyhedral geometry [3], through the fundamental notion of a (Farey) regular simplex [6]. Regular simplexes in $\mathbb{R}^n$ are the affine counterparts of regular cones in $\mathbb{Z}^{n+1}$. Regular fans are complexes of regular cones, and yield a combinatorial classification of nonsingular toric varieties [3].

2. Preliminaries: affine spaces and regular simplexes

A point $y = (y_1, \ldots, y_n) \in \mathbb{R}^n$ is said to be *rational* if each coordinate y_i is a rational number.

Key words and phrases. Affine group over the integers, orbit, $\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$-orbit, complete invariant, rational simplex, Farey regular simplex.

This research was supported by a Marie Curie Intra European Fellowship within the 7th European Community Framework Program (ref. 299401-FP7-PEOPLE-2011-IEF).

2000 *Mathematics Subject Classification.* Primary: 37C85. Secondary: 11B57, 22F05, 37A45, 06F20, 46L80.

A *rational hyperplane* $H \subseteq \mathbb{R}^n$ is a set of the form $H = \{z \in \mathbb{R}^n \mid \langle h, z\rangle = r\}$, for some nonzero vector $h \in \mathbb{Q}^n$ and $r \in \mathbb{Q}$. Here $\langle\text{-},\text{-}\rangle$ denotes scalar product. A *rational affine space* A in $\mathbb{R}^n$ is an intersection of rational hyperplanes in $\mathbb{R}^n$. For any $S \subseteq \mathbb{R}^n$ the *affine hull* $\mathrm{aff}(S)$ is the set of all *affine combinations* in $\mathbb{R}^n$ of elements of S. Thus z belongs to $\mathrm{aff}(S)$ iff there are $w_1, \dots, w_k \in S$ and $\lambda_1, \dots, \lambda_k \in \mathbb{R}$ such that $\lambda_1 + \cdots + \lambda_k = 1$ and $z = \lambda_1 \cdot w_1 + \cdots + \lambda_k \cdot w_k$. A set $\{y_1, \dots, y_m\}$ of points in $\mathbb{R}^n$ is said to be *affinely independent* if none of its elements is an affine combination of the remaining elements. One then easily sees that a subset E of $\mathbb{R}^n$ is a rational affine subspace of $\mathbb{R}^n$ iff there exist $v_0, \dots, v_m \in \mathbb{Q}^n$ such that $E = \mathrm{aff}(v_0, \dots, v_m)$. For $0 \leq m \leq n$, an *m-simplex* in $\mathbb{R}^n$ is the *convex hull* $T = \mathrm{conv}(v_0, \dots, v_m)$ of $m+1$ affinely independent points $v_0, \dots, v_m \in \mathbb{R}^n$. The *vertices* $v_0, \dots, v_m$ are uniquely determined by T. T is said to be a *rational simplex* if its vertices are rational.

By the *denominator* $\mathrm{den}(y)$ of a rational point y we understand the least common denominator of its coordinates. The *homogeneous correspondent* of a rational point $y = (y_1, \dots, y_n) \in \mathbb{R}^n$ is the integer vector

$$\widetilde{y} = (\mathrm{den}(y) \cdot y_1, \dots, \mathrm{den}(y) \cdot y_n,\ \mathrm{den}(y)) \in \mathbb{Z}^{n+1}.$$

Following [6], a rational t-simplex $T = \mathrm{conv}(v_0, \dots, v_t) \subseteq \mathbb{R}^n$ is said to be *(Farey) regular* if the set $\{\tilde{v}_0, \dots, \tilde{v}_t\}$ of homogeneous correspondents of the vertices of T can be extended to a base of the free abelian group $\mathbb{Z}^{n+1}$. (Regular simplexes are called "unimodular" in [5].)

Lemma 1. *For each rational point $x \in \mathbb{R}^n$ and $\psi \in \mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$, $\mathrm{den}(x) = \mathrm{den}(\psi(x))$. Moreover, if $\mathrm{conv}(v_0, \dots, v_n) \subseteq \mathbb{R}^n$ and $\mathrm{conv}(w_0, \dots, w_n) \subseteq \mathbb{R}^n$ are regular n-simplexes, the following conditions are equivalent:*

(i) $\mathrm{den}(v_i) = \mathrm{den}(w_i)$ *for all* $i \in \{0, \dots, n\}$;

(ii) *there exists* $\gamma \in \mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$ *such that* $\gamma(v_i) = w_i$ *for each* $i \in \{0, \dots, n\}$.

Proof. It is easy to see that $\mathrm{den}(x) = \mathrm{den}(\psi(x))$, for every $\psi \in \mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$. So (ii)⇒(i) is immediately verified. For the converse direction (i)⇒(ii), the regularity of $\mathrm{conv}(v_0, \dots, v_n)$ and $\mathrm{conv}(w_0, \dots, w_n)$ yields bases $\{\tilde{v}_0, \dots, \tilde{v}_n\}$ and $\{\tilde{w}_0, \dots, \tilde{w}_n\}$ of the free abelian group $\mathbb{Z}^{n+1}$. There are integer matrices $\mathcal{M}$ and $\mathcal{N}$ such that $\mathcal{M}\tilde{v}_i = \tilde{w}_i$ and $\mathcal{N}\tilde{w}_i = \tilde{v}_i$ for each $i \in \{0, \dots, n\}$. It follows that $\mathcal{M} \cdot \mathcal{N}$ and $\mathcal{N} \cdot \mathcal{M}$ are equal to the identity matrix. For each $i \in \{0, \dots, 1\}$ the last coordinate of $\tilde{v}_i$, as well as the last coordinate of $\tilde{w}_i$, coincide with $\mathrm{den}(v_i) = \mathrm{den}(w_i)$. Thus $\mathcal{M}$ and $\mathcal{N}$ have the form

$$\mathcal{M} = \left(\begin{array}{c|c} \mathcal{U} & b \\ \hline 0, \dots, 0 & 1 \end{array}\right) \quad \mathcal{N} = \left(\begin{array}{c|c} \mathcal{V} & c \\ \hline 0, \dots, 0 & 1 \end{array}\right)$$

for some integer $(n \times n)$-matrices $\mathcal{U}$ and $\mathcal{V}$ and integer vectors $b, c \in \mathbb{Z}^n$. For each $i = 0, \dots, n$ the following holds: $\mathcal{M}\tilde{v}_i = \mathcal{M}(\mathrm{den}(v_i) \cdot (v_i, 1)) = \mathrm{den}(w_i) \cdot (\mathcal{U}v_i + b, 1)$. From $\mathcal{M}\tilde{v}_i = \tilde{w}_i = \mathrm{den}(w_i) \cdot (w_i, 1)$ we obtain $\mathcal{U}v_i + b = w_i$. Similarly, $\mathcal{V}w_i + c = v_i$. In conclusion, both maps $\gamma(x) = \mathcal{U}x + b$ and $\mu(y) = \mathcal{V}y + c$ are in $\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$. Further, $\gamma^{-1} = \mu$ and $\gamma(v_i) = w_i$ for each $i \in \{0, \dots, n\}$. □

Lemma 2. *Let $F \subseteq \mathbb{R}^n$ be an e-dimensional rational affine space ($e = 0, \dots, n$), and v_0 a rational point lying in F. Then there exist rational points $v_1, \dots, v_e \in F$ such that the simplex $\mathrm{conv}(v_0, \dots, v_e)$ is regular.*

Proof. The primitive integer vector $\tilde{v}_0 \in \mathbb{R}^n$ is part of a base $(b_0 = \tilde{v}_0, b_1, \dots, b_n)$ of the free abelian group $\mathbb{Z}^{n+1}$. We may choose the integer vectors $b_1, \dots, b_e$ in such a way that, for each $i = 1, \dots, e$, the uniquely determined rational point v_i in $\mathbb{R}^n$ with $\tilde{v}_i = b_i$ belongs to F. It follows that $\mathrm{conv}(v_0, \dots, v_e)$ is a regular e-simplex contained in F. □

3. Classification of rational affine subspaces of $\mathbb{R}^n$

Given two rational affine spaces $F, G \subseteq \mathbb{R}^n$, the main result of this section (Theorem 8) provides a necessary and sufficient condition for the existence of $\gamma \in \mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$ such that $\gamma(F) = G$. This is a first step for our complete classification of $(\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n)$-orbits in Section 4.

Definition 3. For every non-empty rational affine space $F \subseteq \mathbb{R}^n$ we let

$$\boxed{d_F = \min\{\mathrm{den}(v) \mid v \in F \cap \mathbb{Q}^n\}.}$$

We have the following useful characterization of d_F :

Lemma 4. *Let $F \subseteq \mathbb{R}^n$ be an e-dimensional rational affine space $(e = 0, \ldots, n)$. For any regular e-simplex $\mathrm{conv}(v_0, \ldots, v_e) \subseteq F$, $\quad d_F = \gcd(\mathrm{den}(v_0), \ldots, \mathrm{den}(v_e))$.*

Proof. Let $d = \gcd(\mathrm{den}(v_0), \ldots, \mathrm{den}(v_e))$. For each $v \in F \cap \mathbb{Q}^n$, $\tilde{v} \in \mathbb{Z}\tilde{v}_0 + \cdots + \mathbb{Z}\tilde{v}_e$. Since $\mathrm{den}(v)$ is a linear combination of $\mathrm{den}(v_0), \ldots, \mathrm{den}(v_e)$ with integer coefficients, then $d \leq d_F$. Conversely, for suitable integers $m_0, \ldots, m_e$ we can write $d = m_0 \cdot \mathrm{den}(v_0) + \cdots + m_e \cdot \mathrm{den}(v_e)$. The uniquely determined rational point $w \in F \cap \mathbb{Q}^n$ with $\tilde{w} = m_0 \cdot \tilde{v}_0 + \cdots + m_e \cdot \tilde{v}_e$ satisfies $\mathrm{den}(w) = d$. Thus $d_F \leq d$. □

Lemma 5. *Let $F \subseteq \mathbb{R}^n$ be an e-dimensional rational affine space $(e = 0, \ldots, n)$. Then there are rational points $v_0, \ldots, v_e$ lying in F such that $\mathrm{conv}(v_0, \ldots, v_e)$ is a regular e-simplex and $\mathrm{den}(v_i) = d_F$ for each $i \in \{0, \ldots, e\}$.*

Proof. Pick $v_0 \in F \cap \mathbb{Q}^n$ with $\mathrm{den}(v_0) = d_F$. Lemma 2 yields rational points $w_1, \ldots, w_e \in F$ such that $\mathrm{conv}(v_0, w_1, \ldots, w_e)$ is regular. If $\mathrm{den}(w_i) > d_F$ for some $i \in \{1, \ldots, e\}$, then let the integer m_i be uniquely determined by writing $m_i \cdot d_F < \mathrm{den}(w_i) \leq (m_i + 1) \cdot d_F$. Let $v_i \in F \cap \mathbb{Q}^n$ be the unique rational point with $\tilde{v}_i = \tilde{w}_i - m_i\tilde{v}_0$. Then $d_F \leq \mathrm{den}(v_i) = \mathrm{den}(w_i) - m_i \cdot \mathrm{den}(v_0) = \mathrm{den}(w_i) - m_i \cdot d_F \leq d_F$. The new $(e+1)$-tuple of integer vectors $(\tilde{v}_0, \tilde{v}_1, \ldots, \tilde{v}_e)$ can be completed to a base of $\mathbb{Z}^{n+1}$, precisely as $(\tilde{v}_0, \tilde{w}_1 \ldots, \tilde{w}_e)$ does. Thus $\mathrm{conv}(v_0, \ldots, v_e)$ is regular. □

Definition 6. Let $F \subseteq \mathbb{R}^n$ be an e-dimensional rational affine space. If $0 \leq e < n$ we define

$$\boxed{c_F = \min\{\mathrm{den}(v) \mid v \in \mathbb{Q}^n \setminus F \text{ and } \exists\ v_0, \ldots, v_e \in F \cap \mathbb{Q}^n \text{ with } \mathrm{conv}(v, v_0, \ldots, v_e) \text{ regular}\}.}$$

If $e = n$ we define $c_F = 1$.

Lemma 7. *Let $F \subseteq \mathbb{R}^n$ be an e-dimensional rational affine space $(e = 0, \ldots, n)$. Then there are rational points $v_0, \ldots, v_n$ such that*

(i) $v_0, \ldots, v_e \in F$;
(ii) $\mathrm{den}(v_i) = d_F$ *for all* $i \in \{0, \ldots, e\}$;
(iii) $\mathrm{den}(v_i) = c_F$ *for all* $i \in \{0, \ldots, n\} \setminus \{0, \ldots, e\}$;
(iv) $\mathrm{conv}(v_0, \ldots, v_n)$ *is a regular n-simplex (whence in particular, $v_{e+1}, \ldots, v_n \notin F$).*

Proof. In case $e = n$, the result follows directly from Lemma 5. In case $e < n$, let $w_0, \ldots, w_n \in \mathbb{Q}^n$ be such that $w_0, \ldots, w_e \in F$, $\mathrm{conv}(w_0, \ldots, w_n)$ is regular, and $\mathrm{den}(w_{e+1}) = c_F$. Lemma 5 yields rational points $v_0, \ldots, v_e \in F$ such that $\mathrm{conv}(v_0, \ldots, v_e)$ is regular and $\mathrm{den}(v_i) = d_F$ for each $0 \leq i \leq e$. For every $v \in F \cap \mathbb{Q}^n$ we can write $\tilde{v} \in \mathbb{Z}\tilde{v}_0 + \cdots + \mathbb{Z}\tilde{v}_e$, whence in particular $\mathbb{Z}\tilde{v}_0 + \cdots + \mathbb{Z}\tilde{v}_e = \mathbb{Z}\tilde{w}_0 + \cdots + \mathbb{Z}\tilde{w}_e$. Therefore, $\{\tilde{v}_0, \ldots, \tilde{v}_e, \tilde{w}_{e+1}, \ldots, \tilde{w}_n\}$ is a base of $\mathbb{Z}^{n+1}$, and the n-simplex $T_0 = \mathrm{conv}(v_0, \ldots, v_e, w_{e+1}, \ldots, w_n)$ is regular. In order to obtain from $w_{e+1}, \ldots, w_n$ the points $v_{e+1}, \ldots, v_n$ satisfying (iii)-(iv), we preliminarily set $v_{e+1} = w_{e+1}$. Similarly, for each $i = e+2, \ldots, n$ such that $\mathrm{den}(w_i) = c_F$ we set $v_i = w_i$. If $c_F = \mathrm{den}(w_{e+2}) = \cdots = \mathrm{den}(w_n)$, we are done. Otherwise, for all $i \in \{e+2, \ldots, n\}$ with $\mathrm{den}(w_i) > c_F$, letting the integer m_i be the unique solution of $m_i \cdot c_F < \mathrm{den}(w_i) \leq (m_i + 1) \cdot c_F$, we let the rational point $v_i \in \mathbb{R}^n$ be uniquely determined by the stipulation $\tilde{v}_i = \tilde{w}_i - m_i \cdot \tilde{w}_{e+1}$. We then have $c_F \leq \mathrm{den}(v_i) = \mathrm{den}(w_i) - m_i \cdot \mathrm{den}(w_{e+1}) = \mathrm{den}(w_i) - m_i \cdot c_F \leq c_F$. The n-simplex $T = \mathrm{conv}(v_0, \ldots, v_n)$ obtained from T_0 in this way is regular, and has the desired properties. □

The following theorem yields the classification of $(\mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n)$-orbits of rational affine spaces: the complete classifier is just a triplet of positive integers. As usual, for every affine space $F \subseteq \mathbb{R}^n$, $\dim(F)$ denotes its dimension.

Theorem 8. *Let $F, G \subseteq \mathbb{R}^n$ be non-empty rational affine spaces. Then following conditions are equivalent:*

(i) *There exists* $\gamma \in \mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$ *such that* $\gamma(F) = G$*;*

(ii) $(\dim(F), d_F, c_F) = (\dim(G), d_G, c_G)$.

Proof. (i)⇒(ii) Since $\gamma \in \mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$ is a bijective affine transformation, $\dim(F) = \dim(\gamma(F)) = \dim(G)$. The identities $d_F = d_G$ and $c_F = c_G$ follow directly from Definitions 3 and 6, because γ preserves denominators (Lemma 1).

(ii)⇒(i). Let $v_0, \ldots, v_n \in \mathbb{Q}^n$ satisfy conditions (i)-(iv) of Lemma 7 relative to F. Similarly let $w_0, \ldots, w_n \in \mathbb{Q}^n$ satisfy the same conditions relative to G. By hypothesis, $\mathrm{den}(v_i) = d_F = d_G = \mathrm{den}(w_i)$ for each $0 \leq i \leq \dim(F) = \dim(G)$ and $\mathrm{den}(v_i) = c_F = c_G = \mathrm{den}(w_i)$ for each $i \in \{0, \ldots, n\} \setminus \{0, \ldots, \dim(F)\}$. Lemma 1 yields $\gamma \in \mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$ such that $\gamma(v_i) = w_i$ for each $i \in \{0, \ldots, n\}$. Since γ is an affine transformation and $F = \mathrm{aff}(v_0, \ldots, v_e)$, then $G = \mathrm{aff}(w_0, \ldots, w_e) = \mathrm{aff}(\gamma(v_0), \ldots, \gamma(v_e)) = \gamma(F)$. □

As shown by the next three results, in many cases the parameter c_F has no role in the classification of $(\mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n)$-orbits of rational affine spaces.

Lemma 9. *Let* $F \subseteq \mathbb{R}^n$ *be a rational affine space. If* $0 \leq \dim(F) < n-1$ *then* $c_F = 1$*. If* $\dim(F) = n-1$*, then* $\gcd(c_F, d_F) = 1$ *and* $1 \leq c_F \leq \max(1, d_F/2)$.

Proof. Choose points $v_0, \ldots, v_n \in \mathbb{R}^n$ satisfying conditions (i)-(iv) in Lemma 7. Since $\tilde{v}_0, \ldots, \tilde{v}_n$ is a base of $\mathbb{Z}^{n+1}$, some linear combination of $\mathrm{den}(v_0), \ldots, \mathrm{den}(v_n)$ with integer coefficients equals 1. Thus $\gcd(d_F, c_F) = 1$. A moment's reflection shows that $c_F \leq d_F$. So, in case $d_F = 1$ then $c_F = 1$, and the result follows. On the other hand, in case $d_F > 1$, from $\gcd(d_F, c_F) = 1$ and $c_F \leq d_F$, it follows that $c_F < d_F$.

Let $e = \dim(F)$.

If $e < n-1$, then $v_{e+1} \neq v_n$ and (by Lemma 7(iii)) $\mathrm{den}(v_{e+1}) = \mathrm{den}(v_n) = c_F$. Let $v \in \mathbb{Q}^n$ be such that $\tilde{v} = \tilde{v}_0 - \tilde{v}_n$. Then $\{\tilde{v}_0, \ldots, \tilde{v}_{n-1}, \tilde{v}\}$ is a base of $\mathbb{Z}^{n+1}$. By Definition 6, $c_F \leq \mathrm{den}(v) = d_F - c_F$. Let m be the unique integer such that $m \cdot c_F < \mathrm{den}(v) \leq (m+1) \cdot c_F$. Also let the rational point $w \in \mathbb{R}^n$ be defined by $\tilde{w} = \tilde{v} - m \cdot \tilde{v}_{e+1}$. Observe that $\{\tilde{v}_0, \ldots, \tilde{v}_{n-1}, \tilde{w}\}$ is a base of $\mathbb{Z}^{n+1}$. The minimality property of c_F entails $c_F \leq \mathrm{den}(w) = \mathrm{den}(v) - m \cdot \mathrm{den}(v_{e+1}) = \mathrm{den}(v) - m \cdot c_F \leq c_F$. Then $\mathrm{den}(v) = d_F - c_F = (m+1) \cdot c_F$, whence $d_F = (m+2) \cdot c_F$. Since $\gcd(d_F, c_F) = 1$, then $c_F = 1$.

If $e = n-1$, then $\mathrm{den}(v_n) = c_F$. Let $v \in \mathbb{Q}^n$ be such that $\tilde{v} = \tilde{v}_0 - \tilde{v}_n$. Then $\{\tilde{v}_0, \ldots, \tilde{v}_{n-1}, \tilde{v}\}$ is a base of $\mathbb{Z}^{n+1}$. The minimality property of c_F yields the inequality $c_F \leq \mathrm{den}(v) = d_F - c_F$, that is, $c_F \leq d_F/2$. □

Corollary 10. *Let* $F, G \subseteq \mathbb{R}^n$ *be non-empty rational affine spaces. Suppose* $\dim(F) = \dim(G) \neq n-1$*. Then there exists* $\gamma \in \mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$ *such that* $\gamma(F) = G$ *iff* $d_F = d_G$.

Corollary 11. *Let* $F, G \subseteq \mathbb{R}^n$ *be non-empty rational affine spaces, with* $d_F = d_G \in \{1, 2, 3, 4, 6\}$*. Then there exists* $\gamma \in \mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$ *such that* $\gamma(F) = G$ *iff* $\dim(F) = \dim(G)$.

The following result will find repeated use in the sequel:

Proposition 12. *Let* $n \in \{1, 2, \ldots\}$ *and* e, d, c *be integers such that*

(i) $e \in \{0, \ldots, n-1\}$*,* $d \in \{1, 2, \ldots\}$*, and* $c \in \{1, \ldots, \max(1, d/2)\}$*;*

(ii) $\gcd(d, c) = 1$*;*

(iii) *if* $e \neq n-1$*, then* $c = 1$.

Then for some integer $p \in \{1, \ldots, d\}$ *with* $\gcd(p, d) = 1$*, the rational affine space*

$$F = \{(y_1, \ldots, y_n) \in \mathbb{R}^n \mid y_{e+1} = \cdots = y_n = \tfrac{p}{d}\}$$

satisfies $(\dim(F), d_F, c_F) = (e, d, c)$.

Proof. By (ii), there exist $p \in \{1, \ldots, d\}$ and $q = 0, 1, \ldots$ such that

$$p \cdot c - q \cdot d = 1.$$

The rational affine space $F = \{(y_1, \ldots, y_n) \in \mathbb{R}^n \mid y_{e+1} = \cdots = y_n = \frac{p}{d}\}$ satisfies $\dim(F) = e$ and $\gcd(p, d) = 1$. Let the rational points $v_0, \ldots, v_e \in \mathbb{R}^n$ be defined as follows, for each $i \in \{0, \ldots, e\}$ and $j \in \{1, \ldots, n\}$:

$$(v_i)_j = j\text{th coordinate of } v_i = \begin{cases} \frac{p}{d} & \text{if } j \in \{e+1, \ldots, n\}; \\ 0 & \text{if } j \in \{1, \ldots, e\} \setminus \{i\}; \\ \frac{1}{d} & \text{if } j = i. \end{cases}$$

Thus in particular, $v_0 = (\underbrace{0, \ldots, 0}_{e \text{ times}}, \underbrace{p/d, \ldots, p/d}_{n-e \text{ times}})$. By definition, $\mathrm{den}(v_0) = \cdots = \mathrm{den}(v_e) = d$.

To prove the identities $d_G = d$ and $c_G = c$ we argue by cases.

Case 1: $c = 1$. We then set $p = 1$ and $q = 0$. Let $\xi_1, \ldots, \xi_n$ denote the canonical base of the vector space $\mathbb{R}^n$, and $\mathbf{0} = (0, \ldots, 0)$ the origin of $\mathbb{R}^n$. Then the n-simplex $\mathrm{conv}(v_0, \ldots, v_e, \mathbf{0}, \xi_{e+2}, \ldots, \xi_n)$ is regular. As a matter of fact, the integer $(n+1) \times (n+1)$-matrix $\mathcal{T}$ whose rows are the vectors $\tilde{v}_1, \ldots, \tilde{v}_e, \tilde{v}_0, \tilde{\xi}_{e+2}, \ldots, \tilde{\xi}_n, \tilde{\mathbf{0}}$, is lower triangular with all diagonal elements $= 1$. By Lemma 4, $d_F = \gcd(\mathrm{den}(v_0), \ldots, \mathrm{den}(v_e)) = d$. By Definition 6, $c_F = 1 = c$.

Case 2: $c \neq 1$. By (iii), $e = n - 1$. Let the point $v \in \mathbb{R}^n$ be defined by $v = (0, \ldots, 0, q/c)$. Then $\mathrm{den}(v) = c$. Since $p \cdot c - q \cdot d = 1$, the n-simplex $\mathrm{conv}(v_0, \ldots, v_{n-1}, v)$ is regular. By Lemma 4, $d_F = \gcd(\mathrm{den}(v_0), \ldots, \mathrm{den}(v_{n-1})) = d$. By Definition 6, $c_F \leq c$. There is $w \in \mathbb{Q}^n$ such that $\mathrm{den}(w) = c_F$ and $\mathrm{conv}(v_0, \ldots, v_{n-1}, w)$ is a regular n-simplex. Since $\{\tilde{v}_1, \ldots, \tilde{v}_n, \tilde{w}\}$ is a basis of $\mathbb{Z}^{n+1}$, $\tilde{v}$ is a linear combination of $\{\tilde{v}_1, \ldots, \tilde{v}_n, \tilde{w}\}$ with integer coefficients. Since $p \cdot c - q \cdot d = 1$, we easily see that $c = c_F + k \cdot d$ for some $k \in \mathbb{Z}$. Using now (i) and Lemma 9 we obtain $c_F = c$, which completes the proof. □

4. $(\mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n)$-orbit classification

For each $x = (x_1, \ldots, x_n) \in \mathbb{R}^n$ let

$$\boxed{F_x = \bigcap \{F \subseteq \mathbb{R}^n \mid x \in F \text{ and } F \text{ is a rational affine space}\}.}$$

Also let the group G_x be defined by

$$\boxed{G_x = \mathbb{Z} + x_1\mathbb{Z} + \cdots + x_n\mathbb{Z}.}$$

In other words, G_x is the subgroup of the additive group $\mathbb{R}$ generated by 1 together with the coordinates of x.

Lemma 13. *For each $x \in \mathbb{R}^n$ and $\gamma \in \mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$, $\gamma(F_x) = F_{\gamma(x)}$ and $G_x = G_{\gamma(x)}$.*

Proof. Let $y = \gamma(x)$. Since γ and γ^{-1} preserve rational affine spaces we have the inclusions $F_y \subseteq \gamma(F_x)$ and $F_x \subseteq \gamma^{-1}(F_y)$, whence $F_y \subseteq \gamma(F_x) \subseteq \gamma(\gamma^{-1}(F_y)) = F_y$.

Let $x = (x_1, \ldots, x_n)$ and $y = (y_1, \ldots, y_n)$. Let $\mathcal{U} \in \mathbb{Z}^{n \times n}$ and $b \in \mathbb{Z}^n$ be such that $\gamma(z) = \mathcal{U}z + b$ for each $z \in \mathbb{R}^n$. Since all terms of $\mathcal{U}$ and all coordinates of b are integers, then $y_i \in G_x$ for each $i \in \{1, \ldots, n\}$. Therefore, $G_y \subseteq G_x$. The converse inclusion follows replacing γ by γ^{-1}. □

Lemma 14. *For each $x = (x_1, \ldots, x_n) \in \mathbb{R}^n$, $\mathrm{rank}(G_x) = \dim(F_x) + 1$ and $d_{F_x} = \max\{k \in \mathbb{Z} \mid 1/k \in G_x\}$.*

Proof. Let $e = \dim(F_x)$ and $d = d_{F_x}$. We argue by cases.

Case 1: $e = n$. Then $F_x = \mathbb{R}^n$. Suppose the integers $k_0, \ldots, k_n$ satisfy $k_0 + k_1 \cdot x_1 + \cdots + k_n \cdot x_n = 0$. Letting the vector $v \in \mathbb{Z}^n$ be defined by $v = (k_1, \ldots, k_n)$, we have the identity $\langle x, v \rangle = -k_0$. Since $F_x = \mathbb{R}^n$ then necessarily $k_1 = \cdots = k_n = 0$, and hence $k_0 = \langle x, v \rangle = 0$. We have just proved that $\mathrm{rank}(G_x) = n + 1$, and $G_x \cap \mathbb{Q} = \mathbb{Z}$. It follows that $\max\{k \in \mathbb{Z} \mid 1/k \in G_x\} = 1$. By Definition 3, $d_{F_x} = 1$, which settles the proof of this case.

Case 2: $e < n$. Then Proposition 12 and Theorem 8 jointly yield $p \in \{1, \dots, d\}$ and $\gamma \in \mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$ such that $\gcd(p, d) = 1$ and

$$\gamma(F_x) = \{(y_1, \dots, y_n) \in \mathbb{R}^n \mid y_i = \frac{p}{d} \text{ for each } i = e+1, \dots, n\} = F.$$

Let $\gamma(x) = y = (y_1, \dots, y_e, \frac{p}{d}, \dots, \frac{p}{d})$. By Lemma 13, $G_y = G_x$ and $F_y = \gamma(F_x) = F$. Let $k_0, \dots, k_e$ be such that $k_0 \cdot \frac{1}{d} + k_1 \cdot y_1 + \cdots + k_e \cdot y_e = 0$. Then the rational point $v = (k_1, \dots, k_e, 0, \dots, 0)$ satisfies $\langle v, y \rangle \in \mathbb{Q}$. Since $F_y = F$, we also have $\langle v, z \rangle \in \mathbb{Q}$ for each $z \in F$, whence $v = 0$ and $k_0 \cdot \frac{1}{d} + h_1 \cdot y_1 + \cdots + k_e \cdot y_e = k_0 \cdot \frac{1}{d} = 0$. Having thus proved that $k_i = 0$ for each $i \in \{0, \dots, e\}$, it follows that $\operatorname{rank}(G_x) = \operatorname{rank}(G_y) = e + 1$.

Assume the integer $k > 0$ satisfies $\frac{1}{k} \in G_x = G_y$. Then there exist integers $k_0, \dots, k_e$ satisfying $\frac{1}{k} = k_0 \cdot \frac{1}{d} + k_1 \cdot y_1 + \cdots + k_e \cdot y_e$. It follows that $1 = k \cdot k_0 \cdot \frac{1}{d} + k \cdot k_1 \cdot y_1 + \cdots + k \cdot k_e \cdot y_e$. From $\operatorname{rank}(G_y) = e + 1$ it follows that $k \cdot k_0 = d$. Thus $k \leq d$, and $d = \max\{k \in \mathbb{Z} \mid \frac{1}{k} \in G_x\}$. This settles Case 2 and completes the proof. □

We are now ready to prove the main result of this paper and its corollaries, thus providing a complete classification of $(\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n)$-orbits.

Theorem 15. *For all $x, y \in \mathbb{R}^n$ the following conditions are equivalent:*

(i) *there exists $\gamma \in \mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$ such that $\gamma(x) = y$;*
(ii) $(G_x, c_{F_x}) = (G_y, c_{F_y})$.

Proof. (i)⇒(ii) follows directly from Theorem 8 and Lemma 13. (ii)⇒(i) We argue by cases:

Case 1: $\operatorname{rank}(G_x) = n+1$. By Lemma 14, $\dim(F_x) = \operatorname{rank}(G_x) - 1 = \operatorname{rank}(G_y) - 1 = \dim(F_y) = n$. Since $G_x = G_y$, there are $(n+1) \times (n+1)$ integer matrices $\mathcal{M}$ and $\mathcal{N}$ such that $\mathcal{M}(x_1, \dots, x_n, 1) = (y_1, \dots, y_n, 1)$ and $\mathcal{N}(y_1, \dots, y_n, 1) = (x_1, \dots, x_n, 1)$. Thus $\mathcal{M}$ and $\mathcal{N}$ have the form

$$\mathcal{M} = \left(\begin{array}{c|c} \mathcal{U} & b \\ \hline 0, \dots, 0 & 1 \end{array}\right) \quad \mathcal{N} = \left(\begin{array}{c|c} \mathcal{V} & c \\ \hline 0, \dots, 0 & 1 \end{array}\right)$$

and $\mathcal{N}\mathcal{M}(x_1, \dots, x_n, 1) = (x_1, \dots, x_n, 1)$. Since $\operatorname{rank}(G_x) = n + 1$, $\mathcal{N}\mathcal{M}$ is the identity $(n+1) \times (n+1)$-matrix and $\mathcal{N} = \mathcal{M}^{-1}$. Thus the affine map $\gamma \colon \mathbb{R}^n \to \mathbb{R}^n$ defined by $\gamma(z) = \mathcal{U}z + b$ belongs to $\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$ and $\gamma(x) = y$.

Case 2: $1 \leq \operatorname{rank}(G_x) \leq n$. By Lemma 14, $\dim(F_x) = \operatorname{rank}(G_x) - 1 = \operatorname{rank}(G_y) - 1 = \dim(F_y)$, and $d_{F_x} = \max\{k \in \mathbb{Z} \mid 1/k \in G_x\} = \max\{k \in \mathbb{Z} \mid 1/k \in G_y\} = d_{F_y}$. Let us use the abbreviations

$$e = \dim(F_x) = \dim(F_y), \;\; d = d_{F_x} = d_{F_y}, \;\; c = c_{F_x} = c_{F_y}.$$

Proposition 12 yields an integer p with $1 \leq p \leq d$ such that the affine space

$$F = \{(y_1, \dots, y_n) \in \mathbb{R}^n \mid y_i = \tfrac{p}{d} \text{ for each } i \in \{e+1, \dots, n\}\}$$

satisfies $(\dim(F), d_F, c_F) = (e, d, c)$. By Theorem 8 there are $\gamma_1, \gamma_2 \in \mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$ such that $\gamma_1(F_x) = \gamma_2(F_y) = F$.

Writing for short $x' = \gamma_1(x)$, $y' = \gamma_2(y)$, the proof of (i) amounts to showing

$$y' \in \operatorname{orb}(x'). \tag{1}$$

To this purpose, we define the rational points $v_0, \dots, v_e \in F$ by stipulating that for all $i \in \{0, \dots, e\}$ and $j \in \{1, \dots, n\}$:

$$(v_i)_j = j\text{th coordinate of } v_i = \begin{cases} \frac{p}{d} & \text{if } j \in \{e+1, \dots, n\}; \\ 0 & \text{if } j \in \{1, \dots, e\} \setminus \{i\}; \\ \frac{1}{d} & \text{if } j = i. \end{cases}$$

Then F coincides with $\operatorname{aff}(v_0, \dots, v_e)$, and $\operatorname{conv}(v_0, \dots, v_e)$ is a regular e-simplex. Next let us define the map $\eta \colon F \to \mathbb{R}^e$ by

$$\eta(z_1, \dots, z_n) = (d \cdot z_1, \dots, d \cdot z_e) \text{ for all } z = (z_1, \dots, z_n) \in F.$$

Equivalently, letting $\xi_1, \dots, \xi_e \in \mathbb{R}^e$ denote the vectors of the canonical base of $\mathbb{R}^e$, η is the unique affine transformation such that $\eta(v_0) = \mathbf{0} =$ the origin of $\mathbb{R}^e$, and $\eta(v_i) = \xi_i$ for each $i \in \{1, \dots, e\}$. For any integer m and subgroup G of the additive group $\mathbb{R}$, let us use the self-explanatory notation $m \cdot G = \{m \cdot g \mid g \in G\}$. Then for every $z = (z_1, \dots, z_n) \in F$ we can write

$$G_{\eta(z)} = d \cdot z_1\mathbb{Z} + \cdots + d \cdot z_e\mathbb{Z} + \mathbb{Z} = d \cdot (z_1\mathbb{Z} + \cdots + z_e\mathbb{Z} + \tfrac{1}{d}\mathbb{Z}) = d \cdot G_z. \tag{2}$$

In particular, $G_{\eta(x')} = d \cdot G_{x'} = d \cdot G_x = d \cdot G_y = d \cdot G_{y'} = G_{\eta(y')}$. Since $\operatorname{rank}(G_{\eta(x')}) = e + 1 = \operatorname{rank}(G_{\eta(y')})$, from Lemma 14 we obtain $\dim(F_{\eta(x')}) = e$. The proof of Case 1 above (with e in place of n) yields $\gamma_3 \in \mathsf{GL}(e,\mathbb{Z}) \ltimes \mathbb{Z}^e$ such that

$$\gamma_3(\eta(x')) = \eta(y'). \tag{3}$$

Since η is one-to-one onto $\mathbb{R}^e$, there exist uniquely determined rational points $w_0, \dots, w_e \in F$ such that

$$\eta(w_0) = \gamma_3(\mathbf{0}) \text{ and } \eta(w_i) = \gamma_3(\xi_i) \text{ for each } i \in \{1, \dots, e\}. \tag{4}$$

By (2),

$$\operatorname{den}(w_i) = d \cdot \operatorname{den}(\gamma_3(\xi_i)) = d, \text{ for each } i \in \{1, \dots, e\}. \tag{5}$$

Claim: The e-simplex $\operatorname{conv}(w_0, \dots, w_e) \subseteq F$ is regular.

Indeed, for each $i \in \{1, \dots, e\}$ we have $\widetilde{\eta(w_i)} = \widetilde{\gamma_3(\xi_i)} = (\gamma_3(\xi_i), 1) = (\eta(w_i), 1)$. Thus

$$\{\widetilde{\eta(w_0)}, \dots, \widetilde{\eta(w_e)}\} = \{(\eta(w_0), 1), \dots, (\eta(w_e), 1)\} \text{ is a base of } \mathbb{Z}^e. \tag{6}$$

In case $c = 1$, the proof of Proposition 12 shows that $p = 1$, whence

$$\tilde{w}_i = d \cdot (w_i, 1) = (d \cdot w_i, d) = (\eta(w_i), 1, \dots, 1, d), \text{ for each } i \in \{1, \dots, e\}. \tag{7}$$

Let $\mu_1, \dots, \mu_{n+1}$ denote the canonical base of the free abelian group $\mathbb{Z}^{n+1}$. From (6)-(7) it follows that $\{\tilde{w}_0, \dots, \tilde{w}_e, \mu_{e+2}, \dots, \mu_{n+1}\}$ is a base of $\mathbb{Z}^{n+1}$. Thus the e-simplex $\operatorname{conv}(w_0, \dots, w_e)$ is regular.

In case $c \neq 1$, by Lemma 9, $e = n - 1$. Further, $\tilde{w}_i = d \cdot (w_i, 1) = (d \cdot w_i, d) = (\eta(w_i), p, d)$ for each $i \in \{0, \dots, e\}$. The proof of Proposition 12 provides an integer $q \geq 1$ such that $p \cdot c - q \cdot d = 1$. Upon defining $v = (0, \dots, 0, \frac{q}{c})$, from (6) it follows that the set of integer vectors $\{(\eta(w_0), p, d), \dots, (\eta(w_e), p, d), (0, \dots, 0, q, c)\} = \{\tilde{w}_0, \dots, \tilde{w}_e, \tilde{v}\}$ is a base of $\mathbb{Z}^{n+1}$. Thus, $\operatorname{conv}(w_0, \dots, w_e)$ is regular.

Our claim is settled.

As an immediate consequence of the regularity of the e-simplex $\operatorname{conv}(w_0, \dots, w_e)$, also the n-simplex $\operatorname{conv}(w_0, \dots, w_e, v_{e+1}, \dots, v_n)$ is regular. By (5), $\operatorname{den}(w_i) = \operatorname{den}(v_i) = d$ for each $i \in \{0, \dots, e\}$. Lemma 1 yields $\gamma_4 \in \mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$ such that $\gamma_4(v_i) = w_i$ for $i \in \{0, \dots, e\}$, and $\gamma_4(v_i) = v_i$ for $i \in \{e + 1, \dots, n\}$. It follows that $\gamma_4(F) = F$.

Further,

$$\eta(\gamma_4(z)) = \gamma_3(\eta(z)) \text{ for each } z \in F. \tag{8}$$

As a matter of fact, by (4) we can write $\eta(\gamma_4(v_0)) = \eta(w_0) = \gamma_3(\mathbf{0}) = \gamma_3(\eta(v_0))$ and $\eta(\gamma_4(v_i)) = \eta(w_i) = \gamma_3(\xi_i) = \gamma_3(\eta(v_i))$ for each $i = 1, \dots, n$. Then (8) follows, because η, γ_3 and γ_4 are affine transformations and $F = \operatorname{aff}(v_0, \dots, v_e)$.

Since $\gamma_4(x') \in F$, from (3) and (8) we can write $\eta(\gamma_4(x')) = \gamma_3(\eta(x')) = \eta(y')$. Since the map $\eta\colon F \to \mathbb{R}^e$ is one-to-one, we finally obtain $\gamma_4(x') = y'$, which proves (1) and completes the proof of the theorem. □

Using Lemmas 9 and 14 we obtain the following special cases of Theorem 15:

Corollary 16. *For all $x, y \in \mathbb{R}^n$, if $\operatorname{rank}(G_x) \neq n$, the following conditions are equivalent:*

(i) *There exists $\gamma \in \mathsf{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$, such that $\gamma(x) = y$;*
(ii) *$G_x = G_y$.*

Corollary 17. *For all $x, y \in \mathbb{R}^n$, if $\max\{k \in \mathbb{Z} \mid 1/k \in G_x\} \in \{1, 2, 3, 4, 6\}$, the following conditions are equivalent:*

(i) *There exists $\gamma \in \mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n$, such that $\gamma(x) = y$;*
(ii) *$G_x = G_y$.*

In conclusion, from Lemmas 9 and 14, Proposition 12 and Theorem 15 we obtain:

Corollary 18. *The map $\mathrm{orb}(x) \mapsto (G_x, c_{F_x})$ is a one-one correspondence between $(\mathsf{GL}(n, \mathbb{Z}) \ltimes \mathbb{Z}^n)$-orbits and pairs (G, c), where G is an arbitrary subgroup of the additive group $\mathbb{R}$ having rank $\leq n+1$ and containing 1 among its elements, and c is an integer satisfying the following conditions: if $\mathrm{rank}(G) \neq n$ then $c = 1$; if $\mathrm{rank}(G) = n$, letting $1/d$ be the smallest positive rational in G, then $1 \leq c \leq \max(1, d/2)$ and $\gcd(c, d) = 1$. There are precisely $\max(1, \phi(d)/2)$ possible values for c.*

References

[1] J.S. Dani, Density properties of orbits under discrete groups, *J. Indian Math. Soc.* 39 (1975) 189–218.
[2] A. Guilloux, A brief remark on orbits of SL(2, $\mathbb{Z}$) in the Euclidean plane, *Ergodic Theory Dynam. Systems* 30 (2010) 1101–1109.
[3] G. Ewald, *Combinatorial convexity and algebraic geometry*, Grad. Texts in Math., Vol. 168, Springer-Verlag, New York, 1996.
[4] M. Laurent, A. Nogueira, Approximation to points in the plane by SL(2, $\mathbb{Z}$)-orbits, *J. London Math. Soc.* (2) 85 (2012) 409–429.
[5] D. Mundici, The Haar theorem for lattice-ordered abelian groups with order-unit, *Discrete Contin. Dyn. Syst.* 21 (2008) 537-549.
[6] D. Mundici, Invariant measure under the affine group over $\mathbb{Z}$, *Combin. Probab. Comput.* 23 (2014) 248–268.
[7] D. Mundici, Classifying $(\mathsf{GL}(2, \mathbb{Z}) \ltimes \mathbb{Z}^2)$-orbits by subgroups of $\mathbb{R}$, (preprint available at arXiv:1401.3708)
[8] A. Nogueira, Orbit distribution on $\mathbb{R}^2$ under the natural action of SL(2, $\mathbb{Z}$), *Indag. Math.* (N.S.), 13 (2002) 103–124.
[9] A. Nogueira, Lattice orbit distribution on $\mathbb{R}^2$, *Ergodic Theory Dynam. Systems* 30 (2010) 1201–1214. Erratum, ibid., p. 1215.
[10] E. Witten, SL(2, $\mathbb{Z}$) action on three-dimensional conformal field theories with abelian symmetry, In: *From Fields to Strings: Circumnavigating Theoretical Physics*, Ian Kogan Memorial Collection (in 3 volumes), M. Shifman et al., Eds., World Scientific, Singapore, (2005), pp. 1173–1200.

(L.M. Cabrer) Department of Statistics, Computer Science and Applications, "Giuseppe Parenti", University of Florence, Viale Morgagni 59 – 50134, Florence, Italy
E-mail address: l.cabrer@disia.unifi.it

(D. Mundici) Department of Mathematics and Computer Science "Ulisse Dini", University of Florence, Viale Morgagni 67/A, I-50134 Florence, Italy
E-mail address: mundici@math.unifi.it